arXiv:1606.02431v1 [math.GR] 8 Jun 2016

Finite groups with small number of cyclic subgroups

Wei Zhou

School of Mathematics and Statistics,
Southwest University, Chongqing 400715, P. R. CHINA
zh_great@swu.edu.cn

Abstract

In this note, we study the finite groups with the number of cylic subgroups no greater than 5.

1 Introduction

For a finite group G, let $C(G)$ be the poset of cyclic subgroups of G. Sometimes $C(G)$ can decide the structure of G. For example, $|C(G)| = |G|$ if and only if G is an elementary abelian 2-group. The groups G such that $|G| - |C(G)| \leq 3$ is classified in [2, 3] and [4].

It is well-known that a finite p-group G has eactly one cyclic subgroup of order p if and only if G is cyclic or generalized quaternion group. Hence the groups with small number of cyclic subgroups will be interesting. In this note, we shall study the finite group G with $|C(G)| \leq 5$.

It is esay to see that $|C(G)| = 1$ if and only $G = 1$, and $|C(G)| = 2$ if and only if $G \cong C_p$ for some prime p. In this note, we will focus on the group G such that $3 \leq |C(G)| \leq 5$.

For a finite group G, denote by $\pi_e(G)$ the set of all element orders of G, and by $\pi(G)$ the set of all prime divisors of $|G|$. For any $i \in \pi_e(G)$, denote by $C_i(G)$ the set of all cylic subgroups of order i in G. Throughout this note, let $c_i = |C_i(G)|$.

[0]Support by National Natural Science Foundation of China (Grant No. 11471266).

AMS Subject Classification: 20D15, 20D25

Key words and phrases: finite groups, cyclic subgroups, 2-groups

2 The main result

For a finite group G, we know

$$\begin{aligned} |G| &= \sum_{k\in\pi_e(G)} c_k \cdot \phi(k), \\ |C(G)| &= \sum_{k\in\pi_e(G)} c_k, \end{aligned} \tag{2.1}$$

where ϕ is the Eucler function.

Lemma 2.1 *If $|C(G)| = 3$, then $G \cong C_{p^2}$ for some prime p.*

Proof: We claim that $|\pi(G)| = 1$. Otherwise, by Cauchy theorem, $|\pi(G)| = 2$. Let $|G| = p^a q^b$, where $a, b \geq 1$. By equation 2.1, $c_p = c_q = 1$. Hence there exist $A \lhd G$ and $B \lhd G$ such that $A \cong C_p$ and $B \cong C_q$. Thus $AB \cong C_{pq}$. Note that $|C(C_{pq})| = 4$, a contradiction.

Hence G is a p-groups. Let $|G| = p^n$. By equation 2.1, $c_p = 1$ or 2. If $c_p = 2$, we get that $p^n = 2p + 1$, a contradiction. Hence $c_p = 1$. Then $c_{p^2} = 1$. From equation 2.1, $n = 2$ and $G \cong C_{p^2}$. □

Lemma 2.2 *If $|C(G)| = 4$, then $G \cong C_{pq}, C_{p^3}$.*

Proof: From equation 2.1, we see that $|\pi(G)| \leq 3$. If $\pi(G) = \{p, q, r\}$, then $c_p = c_q = c_r = 1$ and there exist $A, B, C \lhd G$ such that $A \cong C_p$, $B \cong C_q$ and $C \cong C_r$. Thus $ABC \cong C_{pqr}$. Note that $|C(C_{pqr})| > 4$, a contradiction. Now we get the following two cases:

Case 1. $\pi(G) = \{p, q\}$. Let $|G| = p^a q^b$. From equation 2.1, we have that $c_p = 2$ and $c_q = 1$ or $c_p = 1 = c_q$. If $c_p = 2$ and $c_q = 1$, then $p^a q^b = 2p + q - 2$. Since $p^a q^b - 2p - q + 2 \geq pq - 2p - q + 2 = (p - 1)(q - 2)$, we get $q = 2$ and $a = b = 1$. Since $c_2 = 1$, we get $G \cong C_2 \times C_q$, contrary to $c_p = 2$. Therefore $c_p = c_q = 1$, and we can find $A, B \lhd G$ such that $A \cong C_p$ and $B \cong C_q$. Note that $AB \cong C_{pq}$ and $|C(AB)| = 4$. It follows that $G = AB \cong C_{pq}$.

Case 2. $\pi(G) = \{p\}$. Let $|G| = p^n$. We know that $c_p \leq 3$. If $c_p = 3$, then $p^n = 1 + 3(p - 1)$. Hence $p = 2$ and $n = 2$. This is impossible. If $c_p = 2$, then $c_{q^2} = 1$, and $p^n = p^2 + p - 1$. Note that $p^n - p^2 - p + 1 \geq p^3 - p^2 - p + 1 = (p - 1)(p^2 - 1) > 0$. This is impossible. So we get that $c_p = 1$. By [1, Satz 3.8.2], G is a cyclic or $p = 2$ and G is a generalized quaternion group. Clearly, we get $G \cong C_{p^3}$ in this case. □

Lemma 2.3 *If $|C(G)| = 5$, then $G \cong S_3, C_{p^4}, C_3 \times C_3, Q_8$.*

Proof: Let $\pi(G) = \{p_1, \cdots, p_t\}$. By Cauchy theorem, $t \leq 4$. If $t = 4$, then $c_{p_i} = 1$ for $i = 1, \cdots, 4$. Thus G has a normal cyclic subgroup $N \cong C_{p_1p_2p_3p_4}$. We get a contradiction for $|C(C_{p_1p_2p_3p_4})| > 5$. If $t = 3$, from equation 2.1, we get that $c_{p_1} = 1$ or 2, and $c_{p_2} = c_{p_3} = 1$. Let $V, W \lhd G$ such that $V \cong C_{p_2}$ and $W \cong C_{p_3}$. Thus $|C(VW)| = 4$, which implies that $c_{p_1} = 1$. Let $U \lhd G$ such that $U \cong C_{p_1}$. Note that $UVW \cong C_{p_1p_2p_3}$ and $|C(C_{p_1p_2p_3})| > 5$. This is impossible. So we get the following two cases:

Case 1. $t = 2$. Let $|G| = p^aq^b$. Let $P \in Syl_p(G)$ and $Q \in Syl_q(G)$. Suppose that neither P nor Q are cylcic subgroups of prime order. Then $|C(P)| \geq 3$ and $|C(Q)| \geq 3$. Thus $P \cup Q$ contains all the 5 cyclic subgroups of G, and $G = P \cup Q$. This is impossible. So we can assume that $P \cong C_p$, and $|G| = pq^b$.

If $b = 1$, by symmetry, we can assume $P \lhd G$. Note $|C(C_{pq})| = 4$. We get that Q is not normal in G. So $c_q = |G : N_G(Q)| = p$. Then $p = 3$, and $q = 2$. We get $G \cong S_3$.

Now assume $b > 1$. If P is not normal in G, then $c_p = |G : N_G(P)| \geq 1 + p$. We get $p = 2$ and $c_p = 3$. This imples $b = 1$, a contradiction. Hence $P \lhd G$ and $c_p = 1$. Clearly, $|C(Q)| \leq 3$. By lemma 2.1, $Q \cong C_{q^2}$. Now we get $c_{q^2} = |G : N_G(Q)| \geq 3$, contrary to $|C(G)| = 5$.

Case 2. $t = 1$. Then $|G| = p^a$. If $p \geq 5$, then each cyclic subgroup of G must be normal in G for $|C(G)| = 5$, which implies G is a Dedekind group. Thus G is abelian p-groups. Note that $|C_p(C_p \times C_p)| = p + 1 > 5$. It follows that G is cyclic, and then $G \cong C_{p^4}$.

Next assume $p = 3$. Note $|C(C_3 \times C_3)| = 5$. Thus G can not have a proper subgroup isomorphic to $C_3 \times C_3$. If $G \ncong C_3 \times C_3$, then all the subgroup of order 9 are cyclic, which implies G is cyclic by [1, Satz 3.8.4]. So we get $G \cong C_3 \times C_3$ or C_{3^4}.

Finally, we consider the case that $p = 2$. Let $|G| = 2^n$. Since $|C(G)| = 5$, we get $exp(G) = 2^s \leq 2^4$. If $s = 1$, then G is an elementary abelian 2-group of order 2^n and $|C(G)| = 2^n$, a contradicton. If $s = 3$, let $x \in G$ such that $|x| = 2^3$. Since $|C(\langle x \rangle)| = 4$, there is only one cyclic subgroup in $G - \langle x \rangle$. Hence all the elements in $G - \langle x \rangle$ are contained in a cyclic subgroup and have the same order, 2^r say. So we get $|G - \langle x \rangle| = \phi(2^r)$, and $2^n - 8 = 2^{r-1}$, where $r \leq 3$. This is impossible.

Clearly, if $s = 4$, then $G \cong C_{2^4}$. We need only to consider the case that $s = 2$. Let $x \in G$ with $|x| = 4$. Hence we can get exactly two nontrivial cyclic subgroups from $G - \langle x \rangle$. Let 2^r and 2^2 be the order of the two cyclic subgroups, where $r, s \leq 2$. Then $2^n - 4 = 2^{r-1} + 2^{s-1}$. So we get $n = 3$ and $r, s = 2$. Thus $G = Q_8$. □

Combining lemma 2.1, 2.2 and 2.3, we get the following.

Theorem 2.4 *For a finite group G, $|C(G)| \leq 5$ if and only G is a subgroup of C_{p^4} or $G \cong S_3, Q_8, C_3 \times C_3$, or C_{pq}, where p and q are two different primes.*

References

[1] B. Huppert, Endliche Gruppen I, Springer-Verlag, Berlin 1982.

[2] M. Tărnăuceanu, Finite groups with a certain number of cyclic subgroups, Amer. Math. Monthly 122(2015): 275-276.

[3] M. Tărnăuceanu, Finite groups with a certain number of cyclic subgroups II, http://arxiv.org/abs/1604.04974.

[4] W. Zhou, On the number of cyclic subgroups in finite groups, http://arxiv.org/abs/1605.00193.